

# Borel order dimension

Dilip Raghavan  
(joint work with Ming Xiao)

National University of Singapore

Computability and Combinatorics 2023  
University of Connecticut  
Hartford, CT, USA.  
May 21, 2023

# Outline

- 1 Order Dimension
- 2 Borel order dimension
- 3 A dichotomy
- 4 Locally countable orders

# Notation

- $\leq$  is a **quasi order on**  $P$  if  $\leq$  is a reflexive and transitive relation on  $P$ .
- $<$  is a **partial order on**  $P$  if  $<$  is an irreflexive and transitive relation on  $P$ .
- A quasi order  $\leq$  on  $P$  is **linear** or **total** if for any  $x, y \in P$ ,  $x \leq y \vee y \leq x$ .
- A partial order  $<$  on  $P$  is **linear** or **total** if for any  $x, y \in P$ ,  $x < y \vee y < x \vee x = y$ .
- For a quasi order  $\leq$  on  $P$ ,  $E_{\leq}$  is the equivalence relation on  $P$  defined by

$$p E_{\leq} q \iff (p \leq q \wedge q \leq p).$$

- For a quasi order  $\leq$ ,  $x < y$  means  $x \leq y \wedge y \not\leq x$ .  $<$  is a partial order. For a partial order  $<$ ,  $x \leq y$  means  $x < y \vee x = y$ .  $\leq$  is a quasi order with  $E_{\leq} = =$ .

- For a quasi order  $\leq$  on  $P$ ,  $<$  induces a partial order on  $P/E_{\leq}$ , also denoted  $<$ .
- Example 1:  $\mathcal{D} = \langle 2^{\omega}, \leq_T \rangle$ , where  $\leq_T$  is Turing reducibility.
- Example 2:  $\langle \omega^{\omega}, \leq^* \rangle$ , where  $f \leq^* g$  iff  $\forall^{\infty} n \in \omega [f(n) \leq g(n)]$ .

## Definition

A quasi order  $\mathcal{P} = \langle P, \leq \rangle$  is called a **Borel quasi order** if  $P$  is a Polish space and  $\leq$  is a Borel subset of  $P \times P$ .

- $\mathcal{D}$  and  $\langle \omega^\omega, \leq^* \rangle$  are both Borel quasi orders.

## Definition

A quasi order  $\mathcal{P} = \langle P, \leq \rangle$  is said to be **locally countable (locally finite)** if for every  $x \in P$ ,  $\{y \in P : y \leq x\}$  is countable (finite).

- $\mathcal{D}$  is locally countable.
- $\langle \omega^\omega, \leq^* \rangle$  is not locally countable.

## Definition

Suppose  $\leq_0$  and  $\leq$  are both quasi orders on  $P$ .  $\leq$  is said to **extend**  $\leq_0$  if

- 1  $x \leq_0 y \implies x \leq y$  and
- 2  $x E_{\leq_0} y \iff x E_{\leq} y$ ,

for all  $x, y \in P$ .

If  $\leq$  is a linear quasi order which extends  $\leq_0$ , then we say  $\leq$  **linearizes**  $\leq_0$ .

- $\leq$  extends  $\leq_0$  iff
  - (a)  $P/E_{\leq_0} = P/E_{\leq}$  and
  - (b)  $[x] <_0 [y] \implies [x] < [y]$ , for all  $x, y \in P$ .

## Definition

Suppose  $\leq_0$  and  $\leq$  are both quasi orders on  $P$ .  $\leq$  is said to **extend**  $\leq_0$  if

- 1  $x \leq_0 y \implies x \leq y$  and
- 2  $x E_{\leq_0} y \iff x E_{\leq} y$ ,

for all  $x, y \in P$ .

If  $\leq$  is a linear quasi order which extends  $\leq_0$ , then we say  $\leq$  **linearizes**  $\leq_0$ .

- $\leq$  extends  $\leq_0$  iff
  - (a)  $P/E_{\leq_0} = P/E_{\leq}$  and
  - (b)  $[x] <_0 [y] \implies [x] < [y]$ , for all  $x, y \in P$ .
- If  $<$  is a partial order on  $P/E_{\leq_0}$  with  $<_0 \subseteq <$ , then define  $\leq$  on  $P$  by

$$x \leq y \iff \left( x \leq_0 y \vee [x]_{E_{\leq_0}} < [y]_{E_{\leq_0}} \right)$$

- Then  $\leq$  is a quasi order on  $P$  which extends  $\leq_0$  and the partial order induced by  $\leq$  on  $P/E_{\leq_0} = P/E_{\leq}$  is  $<$ .

## Definition (Dushnik–Miller [1], 1941)

For a quasi order  $\mathcal{P} = \langle P, \leq \rangle$ , the **order dimension** (or simply **dimension**) of  $\mathcal{P}$  is the smallest cardinality of a collection of linear orders on  $P/E_{\leq}$  whose intersection is  $<$ .

$\text{odim}(\mathcal{P})$  will denote the order dimension of  $\mathcal{P}$ .

## Fact

The order dimension of  $\mathcal{P}$  is the minimal  $\kappa$  such that  $\langle P/E_{\leq}, < \rangle$  embeds into a product of  $\kappa$  many linear orders (with the coordinate wise ordering on the product).



## Definition (Dushnik–Miller [1], 1941)

For a quasi order  $\mathcal{P} = \langle P, \leq \rangle$ , the **order dimension** (or simply **dimension**) of  $\mathcal{P}$  is the smallest cardinality of a collection of linear orders on  $P/E_{\leq}$  whose intersection is  $<$ .

$\text{odim}(\mathcal{P})$  will denote the order dimension of  $\mathcal{P}$ .

## Fact

The order dimension of  $\mathcal{P}$  is the minimal  $\kappa$  such that  $\langle P/E_{\leq}, < \rangle$  embeds into a product of  $\kappa$  many linear orders (with the coordinate wise ordering on the product).

- $\text{odim}(\mathcal{P})$  is the minimal  $\kappa$  such that there is a sequence  $\langle \leq_i : i \in \kappa \rangle$  of quasi orders on  $P$  extending  $\leq$  such that for any  $x, y \in P$ , if  $x \not\leq y$ , then  $y <_i x$ , for some  $i \in \kappa$ .

# Elementary facts

- The dimension of a linear order is 1.
- The dimension of an antichain is 2.
- The dimension of a (set-theoretic) tree is 2.
- If  $\mathcal{P}$  is an infinite quasi order, then  $\text{odim}(\mathcal{P}) \leq |\mathcal{P}|$ .
- If  $\langle P, \leq \rangle$  embeds into  $\langle Q, \leq_0 \rangle$ , then  $\text{odim}(\langle Q, \leq_0 \rangle) \geq \text{odim}(\langle P, \leq \rangle)$ .

## Locally finite orders

- If  $\mathcal{P}$  is locally finite and  $|P| = \kappa$ , then  $\mathcal{P}$  embeds into  $\langle [\kappa]^{<\aleph_0}, \subseteq \rangle$ .
- So  $\text{odim}(\mathcal{P}) \leq \text{odim}(\langle [\kappa]^{<\aleph_0}, \subseteq \rangle)$ .
- $\text{odim}(\langle [\omega]^{<\aleph_0}, \subseteq \rangle)$  is  $\aleph_0$ .

## Locally finite orders

- If  $\mathcal{P}$  is locally finite and  $|P| = \kappa$ , then  $\mathcal{P}$  embeds into  $\langle [\kappa]^{<\aleph_0}, \subseteq \rangle$ .
- So  $\text{odim}(\mathcal{P}) \leq \text{odim}(\langle [\kappa]^{<\aleph_0}, \subseteq \rangle)$ .
- $\text{odim}(\langle [\omega]^{<\aleph_0}, \subseteq \rangle)$  is  $\aleph_0$ .
- $\text{odim}(\langle [\omega_1]^{<\aleph_0}, \subseteq \rangle)$  is ...

# Locally finite orders

- If  $\mathcal{P}$  is locally finite and  $|P| = \kappa$ , then  $\mathcal{P}$  embeds into  $\langle [\kappa]^{<\aleph_0}, \subseteq \rangle$ .
- So  $\text{odim}(\mathcal{P}) \leq \text{odim}(\langle [\kappa]^{<\aleph_0}, \subseteq \rangle)$ .
- $\text{odim}(\langle [\omega]^{<\aleph_0}, \subseteq \rangle)$  is  $\aleph_0$ .
- $\text{odim}(\langle [\omega_1]^{<\aleph_0}, \subseteq \rangle)$  is  $\dots \aleph_0$ .
- $\text{odim}(\langle [\omega_2]^{<\aleph_0}, \subseteq \rangle)$  is  $\dots$

# Locally finite orders

- If  $\mathcal{P}$  is locally finite and  $|P| = \kappa$ , then  $\mathcal{P}$  embeds into  $\langle [\kappa]^{<\aleph_0}, \subseteq \rangle$ .
- So  $\text{odim}(\mathcal{P}) \leq \text{odim}(\langle [\kappa]^{<\aleph_0}, \subseteq \rangle)$ .
- $\text{odim}(\langle [\omega]^{<\aleph_0}, \subseteq \rangle)$  is  $\aleph_0$ .
- $\text{odim}(\langle [\omega_1]^{<\aleph_0}, \subseteq \rangle)$  is  $\dots \aleph_0$ .
- $\text{odim}(\langle [\omega_2]^{<\aleph_0}, \subseteq \rangle)$  is  $\dots \aleph_0$ .
- $\text{odim}(\langle [\omega_3]^{<\aleph_0}, \subseteq \rangle)$  is  $\dots$

# Locally finite orders

- If  $\mathcal{P}$  is locally finite and  $|P| = \kappa$ , then  $\mathcal{P}$  embeds into  $\langle [\kappa]^{<\aleph_0}, \subseteq \rangle$ .
- So  $\text{odim}(\mathcal{P}) \leq \text{odim}(\langle [\kappa]^{<\aleph_0}, \subseteq \rangle)$ .
- $\text{odim}(\langle [\omega]^{<\aleph_0}, \subseteq \rangle)$  is  $\aleph_0$ .
- $\text{odim}(\langle [\omega_1]^{<\aleph_0}, \subseteq \rangle)$  is  $\dots \aleph_0$ .
- $\text{odim}(\langle [\omega_2]^{<\aleph_0}, \subseteq \rangle)$  is  $\dots \aleph_0$ .
- $\text{odim}(\langle [\omega_3]^{<\aleph_0}, \subseteq \rangle)$  is  $\dots$ 
  - 1 if CH and  $2^{\aleph_1} = \aleph_2$ , then it is  $\aleph_1$ ;
  - 2 else it is  $\aleph_0$ .

## Locally finite orders

- If  $\mathcal{P}$  is locally finite and  $|P| = \kappa$ , then  $\mathcal{P}$  embeds into  $\langle [\kappa]^{<\aleph_0}, \subseteq \rangle$ .
- So  $\text{odim}(\mathcal{P}) \leq \text{odim}(\langle [\kappa]^{<\aleph_0}, \subseteq \rangle)$ .
- $\text{odim}(\langle [\omega]^{<\aleph_0}, \subseteq \rangle)$  is  $\aleph_0$ .
- $\text{odim}(\langle [\omega_1]^{<\aleph_0}, \subseteq \rangle)$  is  $\dots \aleph_0$ .
- $\text{odim}(\langle [\omega_2]^{<\aleph_0}, \subseteq \rangle)$  is  $\dots \aleph_0$ .
- $\text{odim}(\langle [\omega_3]^{<\aleph_0}, \subseteq \rangle)$  is  $\dots$ 
  - 1 if CH and  $2^{\aleph_1} = \aleph_2$ , then it is  $\aleph_1$ ;
  - 2 else it is  $\aleph_0$ .

**Theorem (Kierstead and Milner [5], 1996)**

Let  $\kappa \geq \omega$  be any cardinal. Then  $\text{odim}(\langle [\kappa]^{<\omega}, \subseteq \rangle) = \log_2(\log_2(\kappa))$ .



# Locally countable orders

Theorem (Higuchi, Lempp, R., and Stephan [3], 2019)

*Suppose  $\kappa$  is any cardinal such that  $\text{cf}(\kappa) > \omega$  and  $\mathcal{P} = \langle P, \leq \rangle$  is any locally countable quasi order of size  $\kappa^+$ . Then  $\mathcal{P}$  has dimension at most  $\kappa$ .*

Theorem (Kumar and Raghavan [6], 2020)

*$\mathcal{D} = \langle 2^\omega, \leq_T \rangle$  has the largest order dimension among all locally countable quasi orders of size  $2^{\aleph_0}$ .*

## Theorem (Kumar and Raghavan [6], 2020)

*Each of the following is consistent:*

- 1  $\aleph_1 < \text{odim}(\mathcal{D}) < 2^{\aleph_0}$ ;
- 2  $\text{odim}(\mathcal{D}) = 2^{\aleph_0}$  and  $2^{\aleph_0}$  is weakly inaccessible;
- 3  $\text{odim}(\mathcal{D}) = 2^{\aleph_0} = \aleph_{\omega_1}$ ;
- 4  $\text{odim}(\mathcal{D}) = 2^{\aleph_0} = \aleph_{\omega+1}$ .

- Most Borel quasi orders do not have any Borel linearizations.

Definition (Harrington, Marker, and Shelah [2], 1988)

$\mathcal{P}$  is **thin** if there is no perfect set of pairwise incomparable elements.

Theorem (Harrington, Marker, and Shelah [2], 1988)

If  $\mathcal{P} = \langle P, \leq \rangle$  is a thin Borel quasi order, then for some  $\alpha < \omega_1$ , there is a Borel  $f : P \rightarrow 2^\alpha$  such that

- 1  $x \leq y \implies f(x) \leq_{\text{lex}} f(y)$  and
- 2  $x E_{\leq} y \iff f(x) = f(y)$ , for all  $x, y \in P$ .

- Hence if  $\langle P, \leq_0 \rangle$  is a Borel quasi order and if  $\leq$  is a Borel total quasi order extending  $\leq_0$ , then for some  $\alpha < \omega_1$ , there is a Borel  $f : P \rightarrow 2^\alpha$  such that

$$\begin{aligned}x \leq_0 y &\implies x \leq y \implies f(x) \leq_{\text{lex}} f(y) \text{ and,} \\ x E_{\leq_0} y &\iff x E_{\leq} y \iff f(x) = f(y),\end{aligned}$$

for all  $x, y \in P$ .

- Kanovei [4] found a Borel quasi order  $\langle 2^\omega, \leq_0 \rangle$  which is the canonical obstruction to Borel linearizability.

### Theorem (Kanovei [4], 1998)

Suppose  $\langle P, \leq \rangle$  is a Borel quasi order. Then exactly one of the following two conditions is satisfied:

- 1  $\langle P, \leq \rangle$  is Borel linearizable;
- 2 there is a continuous 1-1 map  $F : 2^\omega \rightarrow P$  such that:
  - (2a)  $a \leq_0 b \implies F(a) \leq F(b)$  and
  - (2b)  $a \not\leq_0 b \implies F(a)$  and  $F(b)$  are  $\leq$ -incomparable.

# Borel order dimension

## Definition

Suppose  $\mathcal{P} = \langle P, \leq \rangle$  is a Borel quasi order. The **Borel order dimension** of  $\mathcal{P}$ , denoted  $\text{odim}_B(\mathcal{P})$ , is the minimal  $\kappa$  such that there is a sequence  $\langle \leq_i : i \in \kappa \rangle$  of Borel quasi orders on  $P$  extending  $\leq$  such that for any  $x, y \in P$ , if  $x \not\leq y$ , then  $y <_i x$ , for some  $i \in \kappa$ .

## Definition

Let  $X$  be a set and  $R$  a binary relation on  $X$  that is disjoint from the diagonal. An  **$R$ -loop** is a finite sequence  $x_0, \dots, x_k \in X$  so that  $(x_i, x_{i+1}) \in R$  for all  $i < k$ ,  $(x_k, x_0) \in R$ .

## Definition

Let  $\mathcal{X} = \langle X, R \rangle$  be a structure as in the previous definition. The **loop-free chromatic number of  $\mathcal{X}$** , denoted  $\mathcal{H}(\mathcal{X})$ , is the minimal  $\kappa$  such that  $X = \bigcup_{\lambda < \kappa} X_\lambda$ , where no  $X_\lambda$  contains an  $R$ -loop.

If  $X$  is a Polish space and  $R$  is a Borel binary relation on  $X$  that is disjoint from the diagonal, then the **Borel loop-free chromatic number of  $\mathcal{X}$** , denoted  $\mathcal{H}_B(\mathcal{X})$ , is the minimal  $\kappa$  such that  $X = \bigcup_{\lambda < \kappa} X_\lambda$ , where each  $X_\lambda$  is a Borel set that does not contain any  $R$ -loops.

- Suppose  $\mathcal{P} = \langle P, \leq \rangle$  is a quasi order. Let  $\mathcal{A}_{\mathcal{P}} = (P \times P) \setminus \geq$  and define  $\mathcal{R}_{\mathcal{P}}$  on  $\mathcal{A}_{\mathcal{P}} \times \mathcal{A}_{\mathcal{P}}$  by  $(p_0, q_0) \mathcal{R}_{\mathcal{P}} (p_1, q_1) \iff q_0 \leq p_1$ .
- $\mathcal{R}_{\mathcal{P}}$  is disjoint from the diagonal because for any  $(p, q) \in \mathcal{A}_{\mathcal{P}}$ ,  $q \not\leq p$ .



- Suppose  $\mathcal{P} = \langle P, \leq \rangle$  is a quasi order. Let  $\mathcal{A}_{\mathcal{P}} = (P \times P) \setminus \geq$  and define  $\mathcal{R}_{\mathcal{P}}$  on  $\mathcal{A}_{\mathcal{P}} \times \mathcal{A}_{\mathcal{P}}$  by  $(p_0, q_0) \mathcal{R}_{\mathcal{P}} (p_1, q_1) \iff q_0 \leq p_1$ .
- $\mathcal{R}_{\mathcal{P}}$  is disjoint from the diagonal because for any  $(p, q) \in \mathcal{A}_{\mathcal{P}}$ ,  $q \not\leq p$ .
- Suppose  $\kappa = \text{odim}(\mathcal{P})$  and that  $\langle \leq_{\lambda} : \lambda < \kappa \rangle$  is a witness.
- Let  $X_{\lambda} = \leq_{\lambda} \setminus \geq$ . Then  $\mathcal{A}_{\mathcal{P}} = \bigcup_{\lambda < \kappa} X_{\lambda}$ .
- If  $(p_0, q_0), \dots, (p_k, q_k)$  is an  $\mathcal{R}_{\mathcal{P}}$ -loop in  $X_{\lambda}$ , then  $p_0 E_{\leq_{\lambda}} q_0$ , which implies  $p_0 E_{\leq} q_0$ , which is impossible as  $q_0 \not\leq p_0$ .
- Hence  $\mathcal{H}(\langle \mathcal{A}_{\mathcal{P}}, \mathcal{R}_{\mathcal{P}} \rangle) \leq \text{odim}(\mathcal{P})$ .

- Conversely suppose  $\mathcal{H}(\langle \mathcal{A}_\mathcal{P}, \mathcal{R}_\mathcal{P} \rangle) = \kappa$  and that  $\langle X_\lambda : \lambda < \kappa \rangle$  is a witness.
- Let  $\leq_\lambda$  be the transitive closure of  $\leq \cup X_\lambda$ .
- $\leq_\lambda$  is then a quasi order on  $P$  and  $\leq \subseteq \leq_\lambda$ .
- $E_{\leq_\lambda} = E_\lambda$  because  $X_\lambda$  is  $\mathcal{R}_\mathcal{P}$ -loop free.

- Conversely suppose  $\mathcal{H}(\langle \mathcal{A}_\mathcal{P}, \mathcal{R}_\mathcal{P} \rangle) = \kappa$  and that  $\langle X_\lambda : \lambda < \kappa \rangle$  is a witness.
- Let  $\leq_\lambda$  be the transitive closure of  $\leq \cup X_\lambda$ .
- $\leq_\lambda$  is then a quasi order on  $P$  and  $\leq \subseteq \leq_\lambda$ .
- $E_{\leq_\lambda} = E_\lambda$  because  $X_\lambda$  is  $\mathcal{R}_\mathcal{P}$ -loop free.
- For example, if  $pX_\lambda qX_\lambda rX_\lambda p$ , then  $(p, q), (q, r), (r, p)$  would be an  $\mathcal{R}_\mathcal{P}$ -loop in  $X_\lambda$ .
- Similarly if  $p \leq qX_\lambda rX_\lambda s \leq tX_\lambda p$ , then  $(q, r), (r, s), (t, p)$  is an  $\mathcal{R}_\mathcal{P}$ -loop in  $X_\lambda$ .

- Conversely suppose  $\mathcal{H}(\langle \mathcal{A}_P, \mathcal{R}_P \rangle) = \kappa$  and that  $\langle X_\lambda : \lambda < \kappa \rangle$  is a witness.
- Let  $\leq_\lambda$  be the transitive closure of  $\leq \cup X_\lambda$ .
- $\leq_\lambda$  is then a quasi order on  $P$  and  $\leq \subseteq \leq_\lambda$ .
- $E_{\leq_\lambda} = E_\lambda$  because  $X_\lambda$  is  $\mathcal{R}_P$ -loop free.
- For example, if  $pX_\lambda qX_\lambda rX_\lambda p$ , then  $(p, q), (q, r), (r, p)$  would be an  $\mathcal{R}_P$ -loop in  $X_\lambda$ .
- Similarly if  $p \leq qX_\lambda rX_\lambda s \leq tX_\lambda p$ , then  $(q, r), (r, s), (t, p)$  is an  $\mathcal{R}_P$ -loop in  $X_\lambda$ .
- If  $q \not\leq p$ , then  $(p, q) \in \mathcal{A}_P = \bigcup_{\lambda < \kappa} X_\lambda$ . So  $p \leq_\lambda q$ , and since  $E_{\leq_\lambda} = E_\leq$ ,  $p <_\lambda q$ .
- Hence  $\text{odim}(\mathcal{P}) \leq \mathcal{H}(\langle \mathcal{A}_P, \mathcal{R}_P \rangle)$
- Conclusion:  $\text{odim}(\mathcal{P}) = \mathcal{H}(\langle \mathcal{A}_P, \mathcal{R}_P \rangle)$ .

## Theorem (R. and Xiao [7])

If  $\mathcal{P}$  is a Borel quasi order, then  $\text{odim}_B(\mathcal{P}) = \mathcal{H}_B(\langle \mathcal{A}_{\mathcal{P}}, \mathcal{R}_{\mathcal{P}} \rangle)$ .

## Theorem (R. and Xiao [7])

If  $\mathcal{P}$  is a Borel quasi order, then  $\text{odim}_B(\mathcal{P}) = \mathcal{H}_B(\langle \mathcal{A}_{\mathcal{P}}, \mathcal{R}_{\mathcal{P}} \rangle)$ .

- Suppose  $s = \langle n_k : k \in \omega \rangle \in \omega^\omega$  is such that  $n_k \geq 2$  and  $n_k \leq n_{k+1}$ , for all  $k \in \omega$ .
- Define  $T(s) = \prod_{k \in \omega} n_k$ .

## Theorem (R. and Xiao [7])

If  $\mathcal{P}$  is a Borel quasi order, then  $\text{odim}_B(\mathcal{P}) = \mathcal{H}_B(\langle \mathcal{A}_{\mathcal{P}}, \mathcal{R}_{\mathcal{P}} \rangle)$ .

- Suppose  $s = \langle n_k : k \in \omega \rangle \in \omega^\omega$  is such that  $n_k \geq 2$  and  $n_k \leq n_{k+1}$ , for all  $k \in \omega$ .
- Define  $T(s) = \prod_{k \in \omega} n_k$ . Let  $D$  be a dense subset of  $T(s)$  that intersects each level exactly once.
- For  $(b_0, b_1) \in [T(s)]$ , define  $(b_0, b_1) \in R_0(D)$  iff there is a  $d \in D$  and an  $x \in \omega^\omega$ , so that either:

$$b_0 = d \frown \langle i \rangle \frown x \text{ and } b_1 = d \frown \langle i+1 \rangle \frown x, \text{ or}$$

$$b_0 = d \frown \langle n_{|d|} - 1 \rangle \frown x \text{ and } b_1 = d \frown \langle 0 \rangle \frown x.$$

- Let  $\mathcal{G}_0(s, D) = \langle [T(s)], R_0(D) \rangle$ .

## Definition

$\mathcal{M} = \{M \subseteq 2^\omega : M \text{ is Borel and meager}\}$ .

$\text{cov}(\mathcal{M}) = \min \{|\mathcal{F}| : \mathcal{F} \subseteq \mathcal{M} \wedge 2^\omega = \bigcup \mathcal{F}\}$ .

## Lemma (R. and Xiao [7])

$\mathcal{H}_B(\mathcal{G}_0(s, D)) \geq \text{cov}(\mathcal{M})$ .

## Proof.

Every Borel non-meager set must contain a loop. □



## Theorem (R. and Xiao [7])

Suppose  $X$  is Polish  $R \subseteq X \times X$  is Borel and disjoint from the diagonal.  
Then either:

- 1  $\mathcal{H}_B(\langle X, R \rangle) \leq \aleph_0$ , or
- 2 there exist  $s, D$ , and a continuous homomorphism  $f : \mathcal{G}_0(s, D) \rightarrow \langle X, R \rangle$ .

## Definition

For  $s$  and  $D$ , define  $\mathcal{P}_0(s, D) = \langle [T(s)] \times 2, \leq_0 \rangle$ , where  $(b_0, i) \leq_0 (b_1, j)$  iff  $i = 0, j = 1$ , and  $(b_0, b_1) \in R_0(D)$ .

- Note that  $\{((b, 1), (b, 0)) : b \in [T(s)]\} \subseteq \mathcal{A}_{\mathcal{P}_0(s, D)}$ .
- Further,  $((b, 1), (b, 0)) \mathcal{R}_{\mathcal{P}_0(s, D)} ((b', 1), (b', 0))$  iff  $(b, 0) \leq_0 (b', 1)$  iff  $b R_0(D) b'$ .
- Therefore, there is a copy of  $\mathcal{G}_0(s, D)$  inside the structure  $\langle \mathcal{A}_{\mathcal{P}_0(s, D)}, \mathcal{R}_{\mathcal{P}_0(s, D)} \rangle$ .
- Hence  $\text{odim}_B(\mathcal{P}_0(s, D)) \geq \text{cov}(\mathcal{M})$ .

## Theorem (R. and Xiao [7])

For any Borel quasi order  $\mathcal{P} = \langle P, \leq \rangle$  exactly one of the following holds:

- 1  $\text{odim}_B(\mathcal{P}) \leq \aleph_0$ .
- 2 There exist  $s, D$ , and a continuous  $f : [T(s)] \times 2 \rightarrow P$  such that:
  - (2a)  $(b_0, 0) \leq_0 (b_1, 1) \implies f((b_0, 0)) \leq f((b_1, 1))$  and
  - (2b) for every  $b \in [T(s)]$ ,  $f((b, 0))$  and  $f((b, 1))$  are  $\leq$ -incomparable.

## Corollary (R. and Xiao [7])

For every Borel quasi order  $\mathcal{P}$ ,  $\text{odim}_B(\mathcal{P})$  is either countable or at least  $\text{cov}(\mathcal{M})$ .

## Theorem (R. and Xiao [7])

For any Borel quasi order  $\mathcal{P} = \langle P, \leq \rangle$  exactly one of the following holds:

- 1  $\text{odim}_B(\mathcal{P}) \leq \aleph_0$ .
- 2 There exist  $s, D$ , and a continuous  $f : [T(s)] \times 2 \rightarrow P$  such that:
  - (2a)  $(b_0, 0) \leq_0 (b_1, 1) \implies f((b_0, 0)) \leq f((b_1, 1))$  and
  - (2b) for every  $b \in [T(s)]$ ,  $f((b, 0))$  and  $f((b, 1))$  are  $\leq$ -incomparable.

## Corollary (R. and Xiao [7])

For every Borel quasi order  $\mathcal{P}$ ,  $\text{odim}_B(\mathcal{P})$  is either countable or at least  $\text{cov}(\mathcal{M})$ .

## Theorem (R. and Xiao [7])

For every Borel quasi order  $\mathcal{P}$ , if  $\text{odim}_B(\mathcal{P})$  is countable, then  $\mathcal{P}$  has a Borel linearization.

# The Turing degrees

- Combining these results with my earlier results with Higuchi, Lempp, and Stephan, we get that  $\text{odim}_B(\mathcal{D})$  is usually strictly bigger than  $\text{odim}(\mathcal{D})$ .
- For example, if  $\text{cf}(\kappa) > \omega$ ,  $2^{\aleph_0} = \kappa^+$ , and  $\text{MA}_\kappa(\text{countable})$  holds. Then  $\text{odim}(\mathcal{D}) \leq \kappa < \kappa^+ = \text{cov}(\mathcal{M}) = \text{odim}_B(\mathcal{D})$ .
- In particular, if PFA holds, then  $\text{odim}(\mathcal{D}) = \aleph_1 < \aleph_2 = \text{odim}_B(\mathcal{D}) = 2^{\aleph_0}$ .

## Theorem (R. and Xiao [7])

*If  $\mathcal{P}$  is a locally finite Borel quasi order, then  $\text{odim}_B(\mathcal{P}) \leq \aleph_0$ .*

## Theorem (R. and Xiao [7])

If  $\mathcal{P}$  is a locally finite Borel quasi order, then  $\text{odim}_B(\mathcal{P}) \leq \aleph_0$ .

- Our dichotomy does not provide any natural upper bound on  $\text{odim}_B(\mathcal{D})$  other than  $2^{\aleph_0}$ .
- So it is natural to wonder whether  $\text{odim}_B(\mathcal{D}) = 2^{\aleph_0}$ .






## Theorem (R. and Xiao [7])

*There is a c.c.c. forcing which forces that for every locally countable Borel quasi order  $\mathcal{P}$ ,  $\text{odim}_B(\mathcal{P}) = \aleph_1$ .*



- So starting with a ground model  $\mathbb{V}$  where  $2^{\aleph_0} = \aleph_{17}$ , there is a cardinal preserving forcing extension in which  $2^{\aleph_0} = \aleph_{17}$  and for every locally countable Borel quasi order  $\mathcal{P}$ ,  $\text{odim}_B(\mathcal{P}) = \aleph_1$ .
- Each  $\mathcal{P}_0(s, D)$  is locally countable. So in this model,  $\mathcal{H}_B(\mathcal{G}_0(s, D)) = \aleph_1 < 2^{\aleph_0}$ , for every  $s$  and  $D$ .



# Bibliography I

-  B. Dushnik and E. W. Miller, *Partially ordered sets*, Amer. J. Math. **63** (1941), 600–610.
-  L. Harrington, D. Marker, and S. Shelah, *Borel orderings*, Trans. Amer. Math. Soc. **310** (1988), no. 1, 293–302.
-  K. Higuchi, S. Lempp, D. Raghavan, and F. Stephan, *On the order dimension of locally countable partial orderings*, Proc. Amer. Math. Soc. **148** (2020), no. 7, 2823–2833.
-  V. Kanovei, *When a partial Borel order is linearizable*, Fund. Math. **155** (1998), no. 3, 301–309.
-  H. A. Kierstead and E. C. Milner, *The dimension of the finite subsets of  $\kappa$* , Order **13** (1996), no. 3, 227–231.

## Bibliography II

-  A. Kumar and D. Raghavan, *Separating families and order dimension of Turing degrees*, Ann. Pure Appl. Logic **172** (2021), no. 5, Paper No. 102911, 19 pp.
-  D. Raghavan and M. Xiao, *Borel order dimension*, preprint (2023).